

3. INVERSE SEMIGROUPS

MARK V. LAWSON

1. INTRODUCTION

Inverse semigroups were introduced in the 1950s by Ehresmann in France, Preston in the UK and Wagner in the Soviet Union as algebraic analogues of pseudogroups of transformations. One of the goals of this article is to give some insight into inverse semigroups by showing that they can in fact be seen as extensions of presheaves of groups by pseudogroups of transformations.

Inverse semigroups can be viewed as generalizations of groups. Group theory is based on the notion of a symmetry; that is, a structure-preserving bijection. Underlying group theory is therefore the notion of a bijection. The set of all bijections from a set X to itself forms a group, $S(X)$, under composition of functions called the symmetric group. Cayley's theorem tells us that each abstract group is isomorphic to a subgroup of a symmetric group. Inverse semigroup theory, on the other hand, is based on the notion of a partial symmetry; that is, a structure-preserving partial bijection. Underlying inverse semigroup theory, therefore, is the notion of a partial bijection (or partial permutation). The set of all partial bijections from X to itself forms a semigroup, $I(X)$, under composition of partial functions called the symmetric inverse monoid. The Wagner-Preston representation theorem tells us that each abstract inverse semigroup is isomorphic to an inverse subsemigroup of a symmetric inverse monoid. However, symmetric inverse monoids and, by extension, inverse semigroups in general, are endowed with extra structure, as we shall see.

The first version of this article was prepared for the *Workshop on semigroups and categories* held at the University of Ottawa between 2nd and 4th May 2010. The second version was prepared for the Field's Institute sponsored workshop *New directions in inverse semigroups* held between 1st and 4th June 2016 again at the University of Ottawa. This version was prepared for my seminar in Bordeaux in April 2018.

2. BASIC DEFINITIONS

In this section, we shall introduce the rudiments of inverse semigroup theory motivated by the properties of the symmetric inverse monoids. Such monoids have not only algebraic structure but also a partial order, a compatibility relation and an underlying groupoid structure all of which can be defined on arbitrary inverse semigroups.

2.1. The theorem of Wagner and Preston. A semigroup S is said to be *inverse* if for each $s \in S$ there exists a unique element s^{-1} such that

$$s = ss^{-1}s \text{ and } s^{-1} = s^{-1}ss^{-1}.$$

Clearly all groups are inverse semigroups.

An *idempotent* in a semigroup is an element e such that $e^2 = e$. Idempotents play an important role in inverse semigroup because the elements $s^{-1}s$ and ss^{-1} are both idempotents. The set of idempotents of S is denoted by $E(S)$. Two special idempotents are the *identity* element, if it exists, and the *zero* element, if it exists. An inverse semigroup with identity is called an *inverse monoid* and an inverse semigroup with zero is called an *inverse semigroup with zero*. An *inverse subsemigroup* of an inverse semigroup is a subsemigroup that is also closed under inverses. If S is an inverse subsemigroup of T and $E(S) = E(T)$ we say that S is a *wide* inverse subsemigroup of T .

The symmetric inverse monoid really is an inverse monoid in the terms of this definition. The only idempotents in $I(X)$ are the identity functions on the subsets of X ; that is, partial functions of the form 1_A where $A \subseteq X$ and 1_A is the identity function on A .

Remark 2.1. The distinction between semigroups and monoids is not a trivial one. A comparison with C^* -algebras will make the point. Commutative C^* -algebras correspond to locally compact spaces whereas the commutative C^* -algebras with identity correspond to compact spaces. See Section 5.

A semigroup S is said to be *regular* if for each $a \in S$ there exists an element b such that $a = aba$ and $b = bab$. The element b is said to be an *inverse* of a . Thus inverse semigroups are the regular semigroups in which each element has a unique inverse. The following result is elementary but fundamental. It was proved independently by Liber in the Soviet Union, and Douglas Munn and Roger Penrose in the UK.¹

Proposition 2.2. *A regular semigroup is inverse if and only if its idempotents commute.*

Proof. Let S be a regular semigroup in which the idempotents commute and let u and v be inverses of x . Then

$$u = uxu = u(xvx)u = (ux)(vx)u,$$

where both ux and vx are idempotents. Thus, since idempotents commute, we have that

$$u = (vx)(ux)u = vxu = (v xv)xu = v(xv)(xu).$$

Again, xv and xu are idempotents and so

$$u = v(xu)(xv) = v(xux)v = vxv = v.$$

Hence $u = v$.

The converse is a little trickier. Observe first that in a regular semigroup the product of two idempotents e and f has an idempotent inverse. To see why, let $x = (ef)'$ be any inverse of ef . Then the element fxe is an idempotent inverse of ef .

Now let S be a semigroup in which every element has a unique inverse. We shall show that $ef = fe$ for any idempotents e and f . By the result above, $f(ef)'e$ is an idempotent inverse of ef . Thus $(ef)' = f(ef)'e$ by uniqueness of inverses, and so $(ef)'$ is an idempotent. Every idempotent is self-inverse, but on the other hand, the inverse of $(ef)'$ is ef . Thus $ef = (ef)'$ by uniqueness of inverses. Hence

¹Allegedly over lunch in St John's College, Cambridge as graduate students.

ef is an idempotent. We have shown that the set of idempotents is closed under multiplication. It follows that fe is also an idempotent. But $ef(fe)ef = (ef)(ef) = ef$, and $fe(ef)fe = fe$ since ef and fe are idempotents. Thus fe and ef are inverses of ef . Hence $ef = fe$. \square

In the symmetric inverse monoid, the product of the idempotents 1_A and 1_B is just $1_{A \cap B}$ and so the commutativity of idempotent multiplication is just a reflection of the fact that the intersection of subsets is commutative.

Inverses in inverse semigroups behave much like inverses in groups.

Lemma 2.3.

- (1) $(s^{-1})^{-1} = s$.
- (2) $(st)^{-1} = t^{-1}s^{-1}$.
- (3) If e is an idempotent then ses^{-1} is an idempotent.

We now characterize the two extreme types of inverse semigroup: those having exactly one idempotent and those consisting of nothing but idempotents.

Proposition 2.4. *All groups are inverse semigroups, and an inverse semigroup is a group if and only if it has a unique idempotent.*

Proof. Clearly, groups are inverse semigroups. Conversely, let S be an inverse semigroup with exactly one idempotent, e say. Then $s^{-1}s = e = ss^{-1}$ for each $s \in S$. But $es = (ss^{-1})s = s = s(s^{-1}s) = se$, and so e is the identity of S . Hence S is a group. \square

Groups are therefore degenerate inverse semigroups.

Recall that a poset is called a (*meet*) *semilattice* if each pair of elements has a greatest lower bound.² The following result leads to the set of idempotents of an inverse semigroup being referred to as its *semilattice of idempotents*.

Proposition 2.5.

- (1) Let S be an inverse semigroup. Then $E(S)$ is a meet semilattice when we define $e \wedge f = ef$.
- (2) All meet semilattices are inverse semigroups, and an inverse in which every element is an idempotent is a meet semilattice.

Proof. (1) Define $e \leq f$ by $e = ef = fe$. Then this is a partial order on $E(S)$, and with respect to this order each pair of idempotents e and f has a greatest lower bound ef .

(2) Let (P, \wedge) be a meet semilattice. Then P is a commutative semigroup in which $e = e \wedge e$ for each element $e \in P$. Thus (P, \wedge) is an inverse semigroup in which every element is idempotent. \square

In the case of the symmetric inverse monoid $I(X)$, result (1) above is just the fact that the semilattice of idempotents of $I(X)$ is isomorphic to the Boolean algebra of all subsets of X .

The following property is often used to show that definitions involving idempotents are self-dual with respect to left and right. It is part of the folklore of the subject but it played an interesting, and rather unexpected role, in Girard's work on linear logic.³

²Usually denoted by \wedge .

³Check out the third bullet-point on page 345 of [5].

Lemma 2.6. *Let S be an inverse semigroup.*

- (1) *For each idempotent e and element s there is an idempotent f such that $es = sf$.*
- (2) *For each idempotent e and element s there is an idempotent f such that $se = fs$.*

Proof. We prove (1) only since the proof of (2) is similar. Put $f = s^{-1}es$ an idempotent. Then $sf = s(s^{-1}es) = (ss^{-1})es = e(ss^{-1})s = es$, using the fact that idempotents commute. \square

Homomorphisms of inverse semigroups are just semigroup homomorphisms. The convention we shall follow is that if S and T are both monoids or both inverse semigroups with zero then their homomorphisms will be required to be monoid homomorphisms or map zeros to zeros, respectively. *Isomorphisms of inverse semigroups* are just semigroup isomorphisms.

Lemma 2.7. *Let $\theta: S \rightarrow T$ be a homomorphism between inverse semigroups.*

- (1) $\theta(s^{-1}) = \theta(s)^{-1}$ for all $s \in S$.
- (2) If e is an idempotent then $\theta(e)$ is an idempotent.
- (3) If $\theta(s)$ is an idempotent then there is an idempotent e in S such that $\theta(s) = \theta(e)$.
- (4) $\text{Im } \theta$ is an inverse subsemigroup of T .
- (5) If U is an inverse subsemigroup of T then $\theta^{-1}(U)$ is an inverse subsemigroup of S .

Proof. (1) Clearly, $\theta(s)\theta(s^{-1})\theta(s) = \theta(s)$ and $\theta(s^{-1})\theta(s)\theta(s^{-1}) = \theta(s^{-1})$. Thus by uniqueness of inverses we have that $\theta(s^{-1}) = \theta(s)^{-1}$.

(2) $\theta(e)^2 = \theta(e)\theta(e) = \theta(e)$.

(3) If $\theta(s)^2 = \theta(s)$, then $\theta(s^{-1}s) = \theta(s^{-1})\theta(s) = \theta(s)^{-1}\theta(s) = \theta(s)^2 = \theta(s)$.

(4) Since θ is a semigroup homomorphism $\text{im } \theta$ is a subsemigroup of T . By (1), $\text{im } \theta$ is closed under inverses.

(5) Straightforward. \square

If $\theta: S \rightarrow T$ is a homomorphism between inverse semigroups then it induces a homomorphism between the semilattices $\mathbf{E}(S)$ and $\mathbf{E}(T)$. If this restricted homomorphism is injective we say that the homomorphism is *idempotent-separating*.

The following result confirms that inverse semigroups are the right abstract counterparts of the symmetric inverse monoids.

Theorem 2.8 (Wagner-Preston representation theorem). *Every inverse semigroup can be embedded in a symmetric inverse monoid.*

Proof. Given an inverse semigroup S we shall construct an injective homomorphism $\theta: S \rightarrow I(S)$. For each element $a \in S$, define $\theta_a: a^{-1}aS \rightarrow aa^{-1}S$ by $\theta_a(x) = ax$. This is well-defined because $aS = aa^{-1}S$ as the following set inclusions show

$$aS = aa^{-1}aS \subseteq aa^{-1}S \subseteq aS.$$

Also $\theta_{a^{-1}}: aa^{-1}S \rightarrow a^{-1}aS$ and $\theta_{a^{-1}}\theta_a$ is the identity on $a^{-1}aS$ and $\theta_a\theta_{a^{-1}}$ is the identity on $aa^{-1}S$. Thus θ_a is a bijection and $\theta_a^{-1} = \theta_{a^{-1}}$. Define $\theta: S \rightarrow I(S)$ by

$\theta(a) = \theta_a$. This is well-defined by the above. Next we show that $\theta_a\theta_b = \theta_{ab}$. If e and f are any idempotents then

$$eS \cap fS = efS.$$

Thus

$$\text{dom } \theta_a \cap \text{im } \theta_b = a^{-1}aS \cap bb^{-1}S = a^{-1}abb^{-1}S.$$

Hence

$$\text{dom}(\theta_a\theta_b) = \theta_b^{-1}(a^{-1}abb^{-1}S) = b^{-1}a^{-1}aS = b^{-1}a^{-1}abS$$

where we use the following subset inclusions

$$b^{-1}a^{-1}aS = b^{-1}bb^{-1}a^{-1}aS = b^{-1}a^{-1}abb^{-1}S \subseteq b^{-1}a^{-1}abS \subseteq b^{-1}a^{-1}aS.$$

Thus $\text{dom}(\theta_a\theta_b) = \text{dom}(\theta_{ab})$. It is immediate from the definitions that $\theta_a\theta_b$ and θ_{ab} have the same effect on elements, and so θ is a homomorphism. It remains to prove that θ is injective. Suppose that $\theta_a = \theta_b$. Then $a = ba^{-1}a$ and $b = ab^{-1}b$ from which $a = b$ readily follows. \square

Example 2.9. Let (X, τ) be a topological space. Consider the collection $I(X, \tau)$ of all homeomorphisms between the open subsets of X . This is not merely a subset of $I(X)$ but also an inverse subsemigroup. It is our first example of what is known as a *pseudogroup of transformations*. Admittedly, in many applications the word ‘pseudogroup’ often implies extra properties that will not concern us here. Pseudogroups of smooth maps between the open subsets of \mathbb{R}^n are used to define differential manifolds. This and similar applications led Ehresmann and Wagner to develop a general theory of pseudogroups with a view to using them in the foundations of differential geometry.

2.2. The natural partial order. In the previous section, we dealt with the algebraic structures on the symmetric inverse monoid: the product and the inverse. But the symmetric inverse monoid $I(X)$ has other structures in addition to its algebraic ones, and these will leave a trace in arbitrary inverse semigroups via the Wagner-Preston representation theorem.

There is a partial ordering on partial bijections called the restriction ordering. Perhaps surprisingly, this order can be characterized algebraically: namely, $f \subseteq g$ if and only if $f = gf^{-1}f$. This motivates our next definition.

On an inverse semigroup, define $s \leq t$ iff $s = ts^{-1}s$.

Lemma 2.10. *The following are equivalent.*

- (1) $s \leq t$.
- (2) $s = te$ for some idempotent e .
- (3) $s = ft$ for some idempotent f .
- (4) $s = ss^{-1}t$.

Proof. (1) \Rightarrow (2). This is immediate.

(2) \Rightarrow (3). This is immediate by Lemma 2.6.

(3) \Rightarrow (4). Suppose that $s = ft$. Then $fs = s$ and so $fss^{-1} = ss^{-1}$. It follows that $s = ss^{-1}t$.

(4) \Rightarrow (1). Suppose that $s = ss^{-1}t$. Then $s = t(t^{-1}ss^{-1}t)$. Put $i = t^{-1}ss^{-1}t$. Then $si = s$ and so $s^{-1}si = s^{-1}s$. It follows that $s = ts^{-1}s$ giving $s \leq t$. \square

We may now establish the main properties of the relation \leq . They are all straightforward to prove in the light of the above lemma.

Proposition 2.11.

- (1) The relation \leq is a partial order.
- (2) If $s \leq t$ then $s^{-1} \leq t^{-1}$.
- (3) If $s_1 \leq t_1$ and $s_2 \leq t_2$ then $s_1 s_2 \leq t_1 t_2$.
- (4) If e and f are idempotents then $e \leq f$ if and only if $e = ef = fe$.
- (5) $s \leq e$ where e is an idempotent implies that e is an idempotent.

Remark 2.12. Property (1) above leads us to dub \leq the *natural partial order* on S . Property (2) needs to be highlighted since readers familiar with lattice-ordered groups might have been expecting something different. Property (3) tells us that the natural partial order is *compatible with the multiplication*. Property (4) tells us that when the natural partial order is restricted to the semilattice of idempotents we get back the usual ordering on the idempotents. Because the natural partial order is defined algebraically it is preserved by homomorphisms.

Our next result tells us that the partial order encodes how far from being a group an inverse semigroup is.

Proposition 2.13. *An inverse semigroup is a group if and only if the natural partial order is the equality relation.*

Proof. Let S be an inverse semigroup in which the natural partial order is equality. If e and f are any two idempotents then $ef \leq e, f$ and so $e = f$. It follows that there is exactly one idempotent and so S is a group by Proposition 2.4. The converse is immediate. \square

In any poset (X, \leq) , a subset $Y \subseteq X$ is said to be an *order ideal* if $x \leq y \in Y$ implies that $x \in Y$. More generally, if Y is any subset of X then define

$$Y^\downarrow = \{x \in X : x \leq y \text{ for some } y \in Y\}.$$

This is the *order ideal generated by Y* . If $y \in X$ then we denote $\{y\}^\downarrow$ by y^\downarrow and call it the *principal order ideal generated by y* .

Property (5) of Proposition 2.11 tells us that the semilattice of idempotents is an order ideal in S with respect to the natural partial order.

Looking below an idempotent we see only idempotents, what happens if we look up? The answer is that we don't necessarily see only idempotents. The symmetric inverse monoid is an example.

Let (X, \leq) be a poset. If Y is any subset of X then define

$$Y^\uparrow = \{x \in X : x \geq y \text{ for some } y \in Y\}.$$

If $Y = \{y\}$ we denote $\{y\}^\uparrow$ by y^\uparrow .

An inverse semigroup S is said to be *E -unitary* if $e \leq s$ where e is an idempotent implies that s is an idempotent. An inverse semigroup with zero S is said to be *E^* -unitary* if $0 \neq e \leq s$ where e is an idempotent implies that s is an idempotent.

Remark 2.14. The reason for having two definitions, depending on whether the inverse semigroup does not or does have a zero, is because an E -unitary inverse semigroup with zero has to be a semilattice since every element is above the zero. Thus the definition of an E -unitary inverse semigroup in the presence of a zero is uninteresting. This bifurcation between inverse semigroups-without-zero and inverse semigroups-with-zero permeates the subject.

2.3. The compatibility relation. As a partially ordered set $I(X)$ has further properties. The meet of any two partial bijections always exists, but joins are a different matter. Given two partial bijections their union is not always another partial bijection; to be so the partial bijections must satisfy a condition that forms the basis of our next definition.

Define $s \sim t$ iff $s^{-1}t, st^{-1} \in E(S)$. This is called the *compatibility relation*. It is reflexive and symmetric but not generally transitive.

Lemma 2.15. *A pair of elements bounded above is compatible.*

Proof. Let $s, t \leq u$. Then $s^{-1}t \leq u^{-1}u$ and $st^{-1} \leq uu^{-1}$ so that $s \sim t$. \square

A subset of an inverse semigroup is said to be *compatible* if the elements are pairwise compatible. If a compatible subset has a least upper bound it is said to have a *join*.

Lemma 2.16. *$s \sim t$ if and only if $s \wedge t$ exists and $\mathbf{d}(s \wedge t) = \mathbf{d}(s) \wedge \mathbf{d}(t)$ and $\mathbf{r}(s \wedge t) = \mathbf{r}(s) \wedge \mathbf{r}(t)$.*

Proof. We prove that st^{-1} is an idempotent if and only if the greatest lower bound $s \wedge t$ of s and t exists and $(s \wedge t)^{-1}(s \wedge t) = s^{-1}st^{-1}t$. The full result then follows by the dual argument. Suppose that st^{-1} is an idempotent. Put $z = st^{-1}t$. Then $z \leq s$ and $z \leq t$, since st^{-1} is an idempotent. Let $w \leq s, t$. Then $w^{-1}w \leq t^{-1}t$ and so $w \leq st^{-1}t = z$. Hence $z = s \wedge t$. Also

$$z^{-1}z = (st^{-1}t)^{-1}(st^{-1}t) = t^{-1}ts^{-1}st^{-1}t = s^{-1}st^{-1}t.$$

Conversely, suppose that $s \wedge t$ exists and $(s \wedge t)^{-1}(s \wedge t) = s^{-1}st^{-1}t$. Put $z = s \wedge t$. Then $z = sz^{-1}z$ and $z = tz^{-1}z$. Thus $sz^{-1}z = tz^{-1}z$, and so $st^{-1}t = ts^{-1}s$. Hence $st^{-1} = ts^{-1}st^{-1}$, which is an idempotent. \square

Since the compatibility relation is not always transitive it is natural to ask when it is. The answer might have been uninteresting but turns out not to be.

Proposition 2.17. *The compatibility relation is transitive if and only if the semigroup is E -unitary.*

Proof. Suppose that \sim is transitive. Let $e \leq s$, where e is an idempotent. Then se^{-1} is an idempotent because $e = se = se^{-1}$, and $s^{-1}e$ is an idempotent because $s^{-1}e \leq s^{-1}s$. Thus $s \sim e$. Clearly $e \sim s^{-1}s$, and so, by our assumption that the compatibility relation is transitive, we have that $s \sim s^{-1}s$. But $s(s^{-1}s)^{-1} = s$, so that s is an idempotent.

Conversely, suppose that S is E -unitary and that $s \sim t$ and $t \sim u$. Clearly $(s^{-1}t)(t^{-1}u)$ is an idempotent and

$$(s^{-1}t)(t^{-1}u) = s^{-1}(tt^{-1})u \leq s^{-1}u.$$

But S is E -unitary and so $s^{-1}u$ is an idempotent. Similarly, su^{-1} is an idempotent. Hence $s \sim u$. \square

An inverse semigroup is said to be a *meet-semigroup* or a \wedge -*semigroup* if it has all binary meets. Unlike the case with joins, there are no preconditions to a pair of elements having a meet.

Proposition 2.18. *An E^* -unitary inverse semigroup is a meet-semigroup.*

Proof. Let s and t be any pair of elements. Suppose that there exists a non-zero element u such that $u \leq s, t$. Then $uu^{-1} \leq st^{-1}$ and uu^{-1} is a non-zero idempotent. Thus st^{-1} is an idempotent. Similarly $s^{-1}t$ is an idempotent. It follows that $s \wedge t$ exists by Lemma 2.16. If the only element below s and t is 0 then $s \wedge t = 0$. \square

In an inverse semigroup with zero there is a refinement of the compatibility relation which is important. Define $s \perp t$ iff $s^{-1}t = 0 = st^{-1}$. This is the *orthogonality relation*. If an orthogonal subset has a least upper bound then it is said to have an *orthogonal join*.

In the symmetric inverse monoid the union of compatible partial bijections is another partial bijection and the union of an orthogonal pair of partial bijections is another partial bijection which is a disjoint union.

Inverse semigroups generalize groups: the single identity of a group is expanded into a semilattice of idempotents. It is possible to go in the opposite direction and contract an inverse semigroup to a group. On an inverse semigroup S define the relation σ by

$$s \sigma t \Leftrightarrow \exists u \leq s, t$$

for all $s, t \in S$.

Theorem 2.19. *Let S be an inverse semigroup.*

- (1) σ is the smallest congruence on S containing the compatibility relation.
- (2) S/σ is a group.
- (3) If ρ is any congruence on S such that S/ρ is a group then $\sigma \subseteq \rho$.

Proof. (1) We begin by showing that σ is an equivalence relation. Reflexivity and symmetry are immediate. To prove transitivity, let $(a, b), (b, c) \in \sigma$. Then there exist elements $u, v \in S$ such that $u \leq a, b$ and $v \leq b, c$. Thus $u, v \leq b$. The set b^\downarrow is a compatible subset and so $u \wedge v$ exists by Lemma 2.15 and Lemma 2.16. But $u \wedge v \leq a, c$ and so $(a, c) \in \sigma$. The fact that σ is a congruence follows from the fact that the natural partial order is compatible with the multiplication. If $s \sim t$ then by Lemma 2.16, the meet $s \wedge t$ exists. Thus $s \sigma t$. It follows that the compatibility relation is contained in the minimum group congruence.

Let ρ be any congruence containing \sim , and let $(a, b) \in \sigma$. Then $z \leq a, b$ for some z . Thus $z \sim a$ and $z \sim b$. By assumption $(z, a), (z, b) \in \rho$. But ρ is an equivalence and so $(a, b) \in \rho$. Thus $\sigma \subseteq \rho$. This shows that σ is the minimum group congruence.

(2) Clearly, all idempotents are contained in a single σ -class (possibly with non-idempotent elements). Consequently, S/σ is an inverse semigroup with a single idempotent. Thus S/σ is a group by Proposition 2.4.

(3) Let ρ be any congruence such that S/ρ is a group. Let $(a, b) \in \sigma$. Then $z \leq a, b$ for some z . Hence $\rho(z) \leq \rho(a), \rho(b)$. But S/ρ is a group and so its natural partial order is equality. Hence $\rho(a) = \rho(b)$. \square

The congruence σ is called the *minimum group congruence* and the group S/σ the *maximum group image* of S . The properties of this congruence lead naturally to the following result on the category of inverse semigroups.

Theorem 2.20. *The category of groups is a reflective subcategory of the category of inverse semigroups.*

Proof. Let S be an inverse semigroup and $\sigma^\natural: S \rightarrow S/\sigma$ the natural homomorphism. Let $\theta: S \rightarrow G$ be a homomorphism to a group G . Then $\ker \theta$ is a group congruence

on S and so $\sigma \subseteq \ker \theta$ by Theorem 2.19. Thus by standard semigroup theory there is a unique homomorphism θ^* from S/σ to G such that $\theta = \theta^*\sigma^{\natural}$. \square

It follows by standard category theory, such as Chapter IV, Section 3 of [22], that there is a functor from the category of inverse semigroups to the category of groups which takes each inverse semigroup S to S/σ . If $\theta: S \rightarrow T$ is a homomorphism of inverse semigroups then the function $\psi: S/\sigma \rightarrow T/\sigma$ defined by $\psi(\sigma(s)) = \sigma(\theta(s))$ is the corresponding group homomorphism (this can be checked directly).

For inverse semigroups with zero the minimum group congruence is not very interesting since the group degenerates to the trivial group. In this case, replacements have to be found.

Remark 2.21. Constructing groups from inverse semigroups might seem a retrograde step but some important groups arise most naturally as maximum group images of inverse semigroups. However, over the past few years it has become apparent that it is the group of units of an inverse monoid that is also of interest. The *group of units* $U(S)$ of the inverse monoid S is defined to be the set of all elements s such that $s^{-1}s = 1 = ss^{-1}$. That is, the elements which are ‘invertible’ in the old-fashioned sense.

2.4. The underlying groupoid. The product we have defined on the symmetric inverse monoid $I(X)$ is not the only one nor perhaps even the most obvious. Given partial bijections f and g we might also want to define fg only when the domain of f is equal to the range of g . When we do this we are regarding f and g as being functions rather than partial functions. With respect to this restricted product $I(X)$ becomes a groupoid. Motivated by this example, define the *restricted product* in an inverse semigroup by $s \cdot t = st$ if $s^{-1}s = tt^{-1}$ and undefined otherwise.

Proposition 2.22. *Every inverse semigroup S is a groupoid with respect to its restricted product.*

Proof. We begin by showing that all idempotents of S are identities of (S, \cdot) . Let $e \in S$ be an idempotent and suppose that $e \cdot x$ is defined. Then $e = xx^{-1}$ and $e \cdot x = ex$. But $ex = (xx^{-1})x = x$. Similarly, if $x \cdot e$ is defined then it is equal to x . We now check that the axioms (C1), (C2) and (C3) hold.

Axiom (C1) holds: suppose that $x \cdot (y \cdot z)$ is defined. Then

$$x^{-1}x = (y \cdot z)(y \cdot z)^{-1} \text{ and } y^{-1}y = zz^{-1}.$$

But

$$(y \cdot z)(y \cdot z)^{-1} = yzz^{-1}y^{-1} = yy^{-1}.$$

Hence $x^{-1}x = yy^{-1}$, and so $x \cdot y$ is defined. Also $(xy)^{-1}(xy) = y^{-1}y = zz^{-1}$. Thus $(x \cdot y) \cdot z$ is defined. It is clear that $x \cdot (y \cdot z)$ is equal to $(x \cdot y) \cdot z$. A similar argument shows that if $(x \cdot y) \cdot z$ exists then $x \cdot (y \cdot z)$ exists and they are equal.

Axiom (C2) holds: suppose that $x \cdot y$ and $y \cdot z$ are defined. We show that $x \cdot (y \cdot z)$ is defined. We have that $x^{-1}x = yy^{-1}$ and $y^{-1}y = zz^{-1}$. Now

$$(yz)(yz)^{-1} = y(zz^{-1})y^{-1} = y(y^{-1}y)y^{-1} = yy^{-1} = x^{-1}x.$$

Thus $x \cdot (y \cdot z)$ is defined. The proof of the converse is straightforward.

Axiom (C3) holds: for each element x we have that $x \cdot (x^{-1}x)$ is defined, and we have seen that idempotents of S are identities. Thus we put $\mathbf{d}(x) = x^{-1}x$. Similarly, we put $xx^{-1} = \mathbf{r}(x)$. It is now clear that (S, \cdot) is a category. The fact that it is a groupoid is immediate. \square

We call (S, \cdot) the *underlying groupoid* of S . The above result leads to the following pictorial representation of the elements of an inverse semigroup. Recall that $\mathbf{d}(s) = s^{-1}s$, which we now call the *domain idempotent* of s , and that $\mathbf{r}(s) = ss^{-1}$, which we now call the *range idempotent* of s . We can regard s as an *arrow*

$$\mathbf{r}(s) \xleftarrow{s} \mathbf{d}(s)$$

In the following result, if you draw a picture and imagine the elements are partial bijections you will see exactly what is going on.

Proposition 2.23. *Let S be an inverse semigroup. Then for any $s, t \in S$ there exist elements s' and t' such that $st = s' \cdot t'$ where the product on the right is the restricted product.*

Proof. Put $e = \mathbf{d}(s)\mathbf{r}(t)$ and define $s' = se$ and $t' = et$. Observe that $\mathbf{d}(s') = e$ and $\mathbf{r}(t') = e$ and that $st = s't'$. \square

At this point, it is natural to define some relations, called *Green's relations*, which can be defined in any semigroup but assume particularly simple forms in inverse semigroups. We define $s\mathcal{L}t$ iff $\mathbf{d}(s) = \mathbf{d}(t)$; $s\mathcal{R}t$ iff $\mathbf{r}(s) = \mathbf{r}(t)$; and $\mathcal{H} = \mathcal{L} \cap \mathcal{R}$ which corresponds to the hom-sets of the underlying groupoid. We define $s\mathcal{D}t$ iff s and t belong to the same connected component of the underlying groupoid. If \mathcal{K} is any one of Green's relation then K_s denotes the \mathcal{K} -class containing s .

Lemma 2.24.

- (1) If $s \leq t$ and either $s\mathcal{L}t$ or $s\mathcal{R}t$ then $s = t$.
- (2) If $s \sim t$ and either $s\mathcal{L}t$ or $s\mathcal{R}t$ then $s = t$.
- (3) If $s \sim t$ and either $\mathbf{d}(s) \leq \mathbf{d}(t)$ or $\mathbf{r}(s) \leq \mathbf{r}(t)$ then $s \leq t$.

Proof. (1) Suppose that $s \leq t$ and $\mathbf{d}(s) = \mathbf{d}(t)$. Then $s = ts^{-1}s = tt^{-1}t = t$.

(2) Suppose that $s \sim t$ and $\mathbf{d}(s) = \mathbf{d}(t)$. Then $s \wedge t$ exists and $\mathbf{d}(s \wedge t) = \mathbf{d}(s)$ by Lemma 2.16. By (1) above $s \wedge t = s$ and $s \wedge t = t$ and so $s = t$.

(3) Suppose that $s \sim t$ and $\mathbf{d}(s) \leq \mathbf{d}(t)$. Then $s \wedge t$ exists and $\mathbf{d}(s \wedge t) = \mathbf{d}(s)$ by Lemma 2.16. Thus $s \wedge t = s$ and so $s \leq t$. \square

If $\theta: S \rightarrow T$ then for each element $s \in S$ the map θ induces a function from L_s to $L_{\theta(s)}$ by restriction. If all these restricted maps are injective (respectively, surjective) we say that θ is *star injective* (respectively, *star surjective*). In the literature, star injective homomorphisms are also referred to as *idempotent-pure* maps on the strength of the following lemma. We shall use this term when referring to congruences.

Lemma 2.25. *Let $\theta: S \rightarrow T$ be a homomorphism between inverse semigroups. The following are equivalent*

- (1) θ is star injective
- (2) Whenever $\theta(s)$ is an idempotent then s is an idempotent.
- (3) The kernel of θ is contained in the compatibility relation.

Proof. (1) \Rightarrow (2). Let θ be star injective and suppose that $\theta(s)$ is an idempotent. Then $\theta(s^{-1}s) = \theta(s)$ since idempotents are self-inverse. But θ is star injective and so $s^{-1}s = s$.

(2) \Rightarrow (3). Let $\theta(s) = \theta(t)$. Then $\theta(s^{-1}s) = \theta(s^{-1}t)$ and so $s^{-1}t$ is an idempotent. By symmetry st^{-1} is an idempotent and so s and t are compatible.

(3) \Rightarrow (1). Let $\theta(s) = \theta(t)$ and $s\mathcal{L}t$. Then $s \sim t$ and so $s = t$ by Lemma 2.24. \square

The E -unitary inverse semigroups also arise naturally in the context of star injective homomorphisms.

Theorem 2.26. *Let S be an inverse semigroup. Then the following conditions are equivalent:*

- (1) S is E -unitary.
- (2) $\sim = \sigma$.
- (3) σ is idempotent pure.
- (4) $\sigma(e) = E(S)$ for any idempotent e .

Proof. (1) \Rightarrow (2). We have already used the fact that the compatibility relation is contained in σ . Let $(a, b) \in \sigma$. Then $z \leq a, b$ for some z . It follows that $z^{-1}z \leq a^{-1}b$ and $zz^{-1} \leq ab^{-1}$. But S is E -unitary and so $a^{-1}b$ and ab^{-1} are both idempotents. Hence $a \sim b$.

(2) \Rightarrow (3). By Lemma 2.25 a congruence is idempotent pure precisely when it is contained in the compatibility relation.

(3) \Rightarrow (4). This is immediate from the definition of an idempotent pure congruence.

(4) \Rightarrow (1) Suppose that $e \leq a$ where e is an idempotent. Then $(e, a) \in \sigma$. But by (4), the element a is an idempotent. \square

The way in which the class of E -unitary inverse semigroups recurs is a reflection of the importance of this class of inverse semigroups in the history of the subject.

In addition to the underlying groupoid, we may sometimes be able to associate another, smaller, groupoid to an inverse semigroup with zero. Let S be an inverse semigroup with zero. An element $s \in S$ is said to be an *atom* if $t \leq s$ implies that $t = 0$ or $t = s$. The set of atoms of S , if non-empty, forms a groupoid called the *minimal groupoid* of S .

Example 2.27. The symmetric inverse monoid $I(X)$ has an interesting minimal groupoid. It consists of those partial bijections whose domains consist of exactly one element of X . This groupoid is isomorphic to the groupoid $X \times X$ with product given by $(x, y)(y, z) = (x, z)$. This is just the groupoid corresponding to the universal relation on X . When X is finite every partial bijection of X can be written as an orthogonal join of elements of the minimal groupoid.

3. SOME EXAMPLES

So far, our range of examples of inverse semigroups is not very extensive. This state of affairs is something we can now rectify using the tools we have available. We describe three examples: groupoids with zero adjoined, presheaves of groups, and semidirect products of semilattices by groups.

3.1. Groupoids with zero adjoined. Category theorists may shudder at this example but a similar idea lies behind the construction of matrix rings from matrix units.

Proposition 3.1. *Groupoids with zero adjoined are precisely the inverse semigroups in which the natural partial order is equality when restricted to the set of non-zero elements.*

Proof. If G is a groupoid then $S = G^0$, the groupoid G with an adjoined zero, is a semigroup when we define all undefined product to be zero. It is an inverse semigroup and the natural partial order is equality when restricted to the non-zero elements.

To prove the converse, let S be an inverse semigroup in which the natural partial order is equality when restricted to the set of non-zero elements. Let s and t be arbitrary elements in S . If $\mathbf{d}(s) = \mathbf{r}(t)$ then st is just the restricted product. Suppose that $\mathbf{d}(s) \neq \mathbf{r}(t)$. Then $\mathbf{d}(s)\mathbf{r}(t) = 0$. It follows that in this case $st = 0$. Thus the only non-zero products in S are the restricted products and the result follows. \square

3.2. Presheaves of groups. The idempotents of an inverse semigroup commute amongst themselves but needn't commute with anything else. The extreme case where they do is interesting. An inverse semigroup is said to be *Clifford* if its idempotents are central. Abelian inverse semigroups are Clifford semigroups and play a central role in the cohomology of inverse semigroups. We show first how to construct examples of Clifford semigroups.

Let (E, \leq) be a meet semilattice, and let $\{G_e : e \in E\}$ be a family of disjoint groups indexed by the elements of E , the identity of G_e being denoted by 1_e . For each pair e, f of elements of E where $e \geq f$ let $\phi_{e,f} : G_e \rightarrow G_f$ be a group homomorphism, such that the following two axioms hold:

(PG1): $\phi_{e,e}$ is the identity homomorphism on G_e .

(PG2): If $e \geq f \geq g$ then $\phi_{f,g}\phi_{e,f} = \phi_{e,g}$.

We call such a family

$$(G_e, \phi_{e,f}) = (\{G_e : e \in E\}, \{\phi_{e,f} : e, f \in E, f \leq e\})$$

a *presheaf of groups (over the semilattice E)*.

Proposition 3.2. *Let $(G_e, \phi_{e,f})$ be a presheaf of groups. Let $S = S(G_e, \phi_{e,f})$ be the union of the G_e equipped with the product defined by:*

$$xy = \phi_{e,e \wedge f}(x)\phi_{f,e \wedge f}(y),$$

where $x \in G_e$ and $y \in G_f$. With respect to this product, S is a Clifford semigroup.

Proof. The product is clearly well-defined. To prove associativity, let $x \in G_e$, $y \in G_f$ and $z \in G_g$ and put $i = e \wedge f \wedge g$. By definition

$$(xy)z = \phi_{e \wedge f, i}(\phi_{e, e \wedge f}(x)\phi_{f, e \wedge f}(y))\phi_{g, i}(z).$$

But

$$\phi_{e \wedge f, i}(\phi_{e, e \wedge f}(x)\phi_{f, e \wedge f}(y)) = \phi_{e \wedge f, i}(\phi_{e, e \wedge f}(x))\phi_{e \wedge f, i}(\phi_{f, e \wedge f}(y)).$$

By axiom (PG2) this simplifies to $\phi_{e, i}(x)\phi_{f, i}(y)$. Thus

$$(xy)z = \phi_{e, i}(x)\phi_{f, i}(y)\phi_{g, i}(z).$$

A similar argument shows that $x(yz)$ likewise reduces to the right-hand side of the above equation. Thus S is a semigroup.

Observe that if $x, y \in G_e$ then xy is just their product in G_e . Thus if $x \in G_e$ and x^{-1} is the inverse of x in the group G_e then

$$x = xx^{-1}x \text{ and } x^{-1} = x^{-1}xx^{-1}$$

by axiom (PG1). Thus S is a regular semigroup.

The idempotents of S are just the identities of the groups G_e , again by axiom (PG1) and $1_e 1_f = 1_{e \wedge f}$. Thus the idempotents commute. We have thus shown that S is an inverse semigroup.

To finish off, let $x \in G_f$. Then

$$1_e x = \varphi_{e, e \wedge f}(1_e) \varphi_{f, e \wedge f}(x) = 1_{e \wedge f} \varphi_{f, e \wedge f}(x) = \varphi_{f, e \wedge f}(x),$$

and similarly, $x 1_e = \varphi_{f, e \wedge f}(x)$. Consequently, the idempotents of S are central. \square

The underlying groupoid of a Clifford semigroup is just a union of groups as the following lemma shows.

Lemma 3.3. *Let S be an inverse semigroup. Then S is Clifford if and only if $s^{-1}s = ss^{-1}$ for every $s \in S$.*

Proof. Let S be a Clifford semigroup and let $s \in S$. Since the idempotents are central $s = s(s^{-1}s) = (s^{-1}s)s$. Thus $ss^{-1} \leq s^{-1}s$. We may similarly show that $s^{-1}s \leq ss^{-1}$, from which we obtain $s^{-1}s = ss^{-1}$.

Suppose now that $s^{-1}s = ss^{-1}$ for all elements s . Let e be any idempotent and s an arbitrary element. Then $(es)^{-1}es = es(es)^{-1}$. That is $s^{-1}es = ss^{-1}e$. Multiplying on the left by s gives $es = se$, as required. \square

We may now characterize Clifford inverse semigroups.

Theorem 3.4. *An inverse semigroup is a Clifford semigroup if and only if it is isomorphic to a presheaf of groups.*

Proof. Let S be a Clifford semigroup. By Lemma 3.3, we know that $s^{-1}s = ss^{-1}$ for all element s . This implies that the underlying groupoid of S is a union of groups. For each idempotent $e \in \mathbf{E}(S)$ define

$$G_e = \{s \in S : \mathbf{d}(s) = e = \mathbf{r}(s)\}.$$

This is a group, the local group at the identity e in the underlying groupoid. By assumption the union of these groups is the whole of S and each element of S belongs to exactly one of these groups. If $e \geq f$ define $\phi_{e,f}: G_e \rightarrow G_f$ by $\phi_{e,f}(a) = af$. This is a well-defined function, because $\mathbf{d}(af) = e$. We show that $(G_e, \phi_{e,f})$ is a presheaf of groups over the semilattice $\mathbf{E}(S)$.

Axiom (PG1) holds: let $e \in \mathbf{E}(S)$ and $a \in G_e$. Then $\phi_{e,e}(a) = ae = aa^{-1}a = a$.

Axiom (PG2) holds: let $e \geq f \geq g$ and $a \in G_e$. Then

$$(\phi_{f,g} \phi_{e,f})(a) = \phi_{f,g}(\phi_{e,f}(a)) = afg = ag = \phi_{e,g}(a).$$

Let T be the inverse semigroup constructed from this presheaf of groups. Let $a \in G_e$ and $b \in G_f$. We calculate their product in this semigroup. By definition

$$\phi_{e,ef}(a) \phi_{f,ef}(b) = aefbef = afbe = aefb = ab.$$

Thus S and T are isomorphic.

The converse was proved in Proposition 3.2. \square

3.3. Semidirect products of semilattices by groups. The group G acts on the set Y (on the left) if there is a function $G \times Y \rightarrow Y$ denoted by $(g, e) \mapsto g \cdot e$ satisfying $1 \cdot e = e$ for all $e \in Y$ and $g \cdot (h \cdot e) = (gh) \cdot e$ for all $g, h \in G$ and $e \in Y$. If Y is a partially ordered set, then we say that G acts on Y by *order automorphisms* if for all $e, f \in Y$ we have that

$$e \leq f \Leftrightarrow g \cdot e \leq g \cdot f.$$

Observe that in the case of a group action, it is enough to assume that $e \leq f$ implies $g \cdot e \leq g \cdot f$, because if $g \cdot e \leq g \cdot f$ then $g^{-1} \cdot (g \cdot e) \leq g^{-1} \cdot (g \cdot f)$ and so $1 \cdot e \leq 1 \cdot f$, which gives $e \leq f$. If Y is a meet semilattice on which G acts by order automorphisms, then it is automatic that

$$g \cdot (e \wedge f) = g \cdot e \wedge g \cdot f$$

for all $g \in G$ and $e, f \in Y$.

Let $P(G, Y)$ be the set $Y \times G$ equipped with the multiplication

$$(e, g)(f, h) = (e \wedge g \cdot f, gh).$$

Proposition 3.5. *$P(G, Y)$ is an E -unitary inverse semigroup in which the semilattice of idempotents is isomorphic to (Y, \leq) and G is isomorphic to the maximum group homomorphic image of $P(G, Y)$.*

Proof. $P(G, Y)$ is an inverse semigroup in which the inverse of (e, g) is the element $(g^{-1} \cdot e, g^{-1})$, and the idempotents of $P(G, Y)$ are the elements of the form $(e, 1)$. From the definition of the multiplication in $P(G, Y)$ the function $(e, 1) \mapsto e$ is an isomorphism of semilattices. The natural partial order is given by

$$(e, g) \leq (f, h) \Leftrightarrow e \leq f \text{ and } g = h.$$

If $(e, 1) \leq (f, g)$ then $g = 1$ and so $P(G, Y)$ is E -unitary. It also follows from the description of the natural partial order that $(e, g) \sigma(f, h)$ if and only if $g = h$. \square

We may now characterize those inverse semigroups isomorphic to semidirect products of semilattices by groups using many of the ideas introduced in Section 2 to do so.

Theorem 3.6. *Let S be an inverse semigroup. Then the following are equivalent:*

- (1) *The semigroup S is isomorphic to a semidirect product of a semilattice by a group.*
- (2) *S is E -unitary and for each $a \in S$ and $e \in \mathbf{E}(S)$ there exists $b \in S$ such that $b \sim a$ and $b^{-1}b = e$.*
- (3) *$\sigma^{\natural}: S \rightarrow S/\sigma$ is star bijective.*
- (4) *There is a star bijective homomorphism from S to a group.*
- (5) *The function $\theta: S \rightarrow \mathbf{E}(S) \times S/\sigma$ defined by $\theta(a) = (a^{-1}a, \sigma(a))$ is a bijection.*
- (6) *The function $\phi: S \rightarrow \mathbf{E}(S) \times S/\sigma$ defined by $\phi(a) = (aa^{-1}, \sigma(a))$ is a bijection.*

Proof. (1) \Rightarrow (2). Without loss of generality, we may assume that S is a semidirect product of a meet semilattice Y by a group G . The semigroup S is E -unitary by Theorem 3.6. Let $(e, g) \in S$ and $(f, 1) \in \mathbf{E}(S)$. Then the element $(g \cdot f, g)$ of S satisfies

$$(g \cdot f, g) \sim (e, g) \text{ and } (g \cdot f, g)^{-1}(g \cdot f, g) = (f, 1)$$

as required.

(2) \Rightarrow (3). Since S is E -unitary, the homomorphism $\sigma^{\natural}: S \rightarrow S/\sigma$ is star injective by Theorem 2.26. Let $e \in \mathbf{E}(S)$ and $\sigma(a) \in S/\sigma$. By assumption there exists $b \in S$ such that $b^{-1}b = e$ and $b \sim a$. But $b \sim a$ implies $\sigma(b) = \sigma(a)$. Thus σ^{\natural} is also star surjective.

(3) \Rightarrow (4). Immediate.

(4) \Rightarrow (3). Let $\theta: S \rightarrow G$ be a star bijective homomorphism to a group G . Since σ is the minimum group congruence, $\sigma \subseteq \ker \theta$ by Theorem 2.19. But θ is star injective by assumption, and so σ^{\natural} is idempotent pure by Lemma 2.25. In particular, S is E -unitary by Theorem 2.26.

To show that σ^{\natural} is star surjective, let $s \in S$ and $e \in \mathbf{E}(S)$. There exists $t \in S$ such that $t^{-1}t = e$ and $\theta(t) = \theta(s)$, since θ is star surjective. Now $\theta(s^{-1}t)$ is the identity of G , and so $s^{-1}t$ is an idempotent of S since θ is star injective. Similarly, st^{-1} is an idempotent. Hence $s \sim t$ and so $(s, t) \in \sigma$. Thus for each $e \in \mathbf{E}(S)$ and $\sigma(s) \in S/\sigma$, there exists $t \in S$ such that $t^{-1}t = e$ and $\sigma(t) = \sigma(s)$. Thus σ^{\natural} is star surjective.

(3) \Rightarrow (5). Straightforward.

(5) \Rightarrow (6). Suppose that $\phi(a) = \phi(b)$. Then $aa^{-1} = bb^{-1}$ and $\sigma(a) = \sigma(b)$. But $\sigma(a^{-1}) = \sigma(b^{-1})$ and so $\theta(a^{-1}) = \theta(b^{-1})$. By assumption θ is bijective and so $a^{-1} = b^{-1}$, giving $a = b$. Hence ϕ is injective.

Now let $(e, \sigma(s)) \in \mathbf{E}(S) \times S/\sigma$. Since θ is surjective there exists $t \in S$ such that $\theta(t) = (e, \sigma(s^{-1}))$. Thus $t^{-1}t = e$ and $t\sigma s^{-1}$. Hence t^{-1} is such that $t^{-1}\sigma s$ and $t^{-1}(t^{-1})^{-1} = e$. Thus $\phi(t^{-1}) = (e, \sigma(s))$, and so ϕ is surjective.

(6) \Rightarrow (5). A similar argument to (5) \Rightarrow (6).

(6) \Rightarrow (1). We shall use the fact that both the functions ϕ and θ defined above are bijections.

First of all S is E -unitary. For suppose that $e \leq a$ where e is an idempotent. Then $\sigma(e) = \sigma(a)$, and $\sigma(e) = \theta(a^{-1}a)$, so that $\sigma(a) = \sigma(a^{-1}a)$. Thus $\theta(a) = \theta(a^{-1}a)$, and so $a = a^{-1}a$, since θ is a bijection.

We shall define an action of S/σ on $\mathbf{E}(S)$ using θ , and then show that ϕ defines an isomorphism from the semidirect product of $\mathbf{E}(S)$ by S/σ to S .

Define $\sigma(s) \cdot e = tt^{-1}$ where $\theta(t) = (e, \sigma(s))$. This is well-defined because θ is a bijection. The two defining properties of an action hold. Firstly, if $\sigma(e)$ is the identity of S/σ then $\theta(e) = (e, \sigma(e))$ and so $\sigma(e) \cdot e = e$; secondly, $\sigma(u) \cdot (\sigma(v) \cdot e) = \sigma(u) \cdot aa^{-1}$ where $\theta(a) = (e, \sigma(v))$, and $\sigma(u) \cdot aa^{-1} = bb^{-1}$ where $\theta(b) = (aa^{-1}, \sigma(u))$. Now $a\sigma v$ and $b\sigma u$ so that $ba\sigma uv$. Also $a^{-1}a = e$ and $b^{-1}b = aa^{-1}$ so that $(ba)^{-1}ba = a^{-1}a$. Hence $\theta(ba) = (e, \sigma(uv))$. Thus

$$\sigma(uv) \cdot e = (ba)(ba)^{-1} = bb^{-1} = \sigma(u) \cdot (\sigma(v) \cdot e).$$

Next, we show that S/σ acts on $\mathbf{E}(S)$ by means of order automorphisms. Suppose that $e \leq f$. Then $\sigma(a) \cdot e = uu^{-1}$ and $\sigma(a) \cdot f = vv^{-1}$ where

$$\theta(u) = (e, \sigma(a)) \text{ and } \theta(v) = (f, \sigma(a)).$$

Consequently, $e = u^{-1}u$ and $f = v^{-1}v$ and $u\sigma v$. But S is E -unitary, and so σ is equal to the compatibility relation by Theorem 2.26. From $u^{-1}u \leq v^{-1}v$ and $u \sim v$ we obtain $u \leq v$ by Lemma 2.24. Hence $uu^{-1} \leq vv^{-1}$ and so $\sigma(a) \cdot e \leq \sigma(a) \cdot f$.

It only remains to prove that ϕ is a homomorphism. By definition

$$\phi(a)\phi(b) = (aa^{-1}, \sigma(a))(bb^{-1}, \sigma(b)) = (aa^{-1} \wedge \sigma(a) \cdot bb^{-1}, \sigma(ab)).$$

But $\sigma(a) \cdot bb^{-1} = tt^{-1}$ where $\theta(t) = (bb^{-1}, \sigma(a))$. Thus

$$\phi(a)\phi(b) = (aa^{-1}tt^{-1}, \sigma(ab))$$

whereas

$$\phi(ab) = (ab(ab)^{-1}, \sigma(ab)).$$

It remains to show that $aa^{-1}tt^{-1} = ab(ab)^{-1}$. We know that $t^{-1}t = bb^{-1}$ and $t\sigma a$. But $t \sim a$ since S is E -unitary. Thus $tt^{-1}a = at^{-1}t = abb^{-1}$ by Lemma 2.16. Hence $tt^{-1}aa^{-1} = abb^{-1}a^{-1} = ab(ab)^{-1}$. \square

4. FUNDAMENTAL INVERSE SEMIGROUPS

The examples in the last section can be viewed as showing that various natural ways of combining groups and semilattices lead to interesting classes of inverse semigroups. But what does the ‘generic’ inverse semigroup look like? The main goal of this section is to justify the claim made in the Introduction that inverse semigroups should be viewed as common generalizations of presheaves of groups and pseudogroups of transformations. We shall also characterize the congruence-free inverse semigroups with zero.

4.1. The Munn representation. The symmetric inverse monoid is constructed from an arbitrary set. We now show how to construct an inverse semigroup from a meet semilattice. Let (E, \leq) be a meet semilattice, and denote by T_E be the set of all order isomorphisms between principal order ideals of E . Clearly, T_E is a subset of $I(E)$. In fact we have the following.

Proposition 4.1. *The set T_E is an inverse subsemigroup of $I(E)$ whose semilattice of idempotents is isomorphic to E .*

T_E is called the *Munn semigroup* of the semilattice E .

Theorem 4.2 (Munn representation theorem). *Let S be an inverse semigroup. Then there is an idempotent-separating homomorphism $\delta: S \rightarrow T_{E(S)}$ whose image is a wide inverse subsemigroup of $T_{E(S)}$.*

Proof. For each $s \in S$ define the function

$$\delta_s: (s^{-1}s)^\downarrow \rightarrow (ss^{-1})^\downarrow$$

by $\delta_s(e) = ses^{-1}$. We first show that δ_s is well-defined. Let $e \leq s^{-1}s$. Then $ss^{-1}\delta_s(e) = \delta_s(e)$, and so $\delta_s(e) \leq ss^{-1}$. To show that δ_s is order-preserving, let $e \leq f \in (s^{-1}s)^\downarrow$. Then

$$\delta_s(e)\delta_s(f) = ses^{-1}sf s^{-1} = sefs^{-1} = \delta_s(e).$$

Thus $\delta_s(e) \leq \delta_s(f)$.

Consider now the function $\delta_{s^{-1}}: (ss^{-1})^\downarrow \rightarrow (s^{-1}s)^\downarrow$. This is order-preserving by the argument above. For each $e \in (s^{-1}s)^\downarrow$, we have that

$$\delta_{s^{-1}}(\delta_s(e)) = \delta_{s^{-1}}(ses^{-1}) = s^{-1}ses^{-1}s = e.$$

Similarly, $\delta_s(\delta_{s^{-1}}(f)) = f$ for each $f \in (ss^{-1})^\downarrow$. Thus δ_s and $\delta_{s^{-1}}$ are mutually inverse, and so δ_s is an order isomorphism.

Define $\delta: S \rightarrow T_{E(S)}$ by $\delta(s) = \delta_s$. To show that δ is a homomorphism, we begin by calculating $\text{dom}(\delta_s\delta_t)$ for any $s, t \in S$. We have that

$$\text{dom}(\delta_s\delta_t) = \delta_t^{-1}((s^{-1}s)^\downarrow) \cap (tt^{-1})^\downarrow = \delta_t^{-1}((s^{-1}stt^{-1})^\downarrow).$$

But $\delta_t^{-1} = \delta_{t^{-1}}$ and so

$$\text{dom}(\delta_s\delta_t) = ((st)^{-1}st)^\downarrow = \text{dom}(\delta_{st}).$$

If $e \in \text{dom} \delta_{st}$ then

$$\delta_{st}(e) = (st)e(st)^{-1} = s(tet^{-1})s^{-1} = \delta_s(\delta_t(e)).$$

Hence $\delta_s \delta_t = \delta_{st}$.

To show that δ is idempotent-separating, suppose that $\delta(e) = \delta(f)$ where e and f are idempotents of S . Then $\text{dom } \delta(e) = \text{dom } \delta(f)$. Thus $e = f$.

The image of δ is a wide inverse subsemigroup of $T_{\mathbf{E}(S)}$ because every idempotent in $T_{\mathbf{E}(S)}$ is of the form $1_{[e]}$ for some $e \in \mathbf{E}(S)$, and $\delta_e = 1_{[e]}$. \square

The Munn representation should be contrasted with the Wagner-Preston representation: that was injective whereas this has a non-trivial kernel which we shall now describe. The kernel of δ is the congruence μ defined by $(s, t) \in \mu$ if and only if $\mathbf{d}(s) = \mathbf{d}(t)$, $\mathbf{r}(s) = \mathbf{r}(t)$ and for all idempotents e such that $e \leq s^{-1}s$ we have that $ses^{-1} = tet^{-1}$. The definition can be slightly weakened.

Lemma 4.3. *The congruence μ is defined by*

$$(s, t) \in \mu \Leftrightarrow (\forall e \in \mathbf{E}(S)) ses^{-1} = tet^{-1}.$$

Proof. Define $(s, t) \in \mu'$ iff $ses^{-1} = tet^{-1}$ for all idempotents e . We shall prove that $\mu = \mu'$. Observe first that μ' is a congruence. It is clearly an equivalence relation. Suppose that $(a, b) \in \mu'$ and $(c, d) \in \mu'$. The proof that $(ac, bd) \in \mu'$ is straightforward. It follows that from $(s, t) \in \mu'$ we may deduce that $(s^{-1}, t^{-1}) \in \mu'$. Let $(s, t) \in \mu'$. We prove that $(s, t) \in \mu$. To do this we need to prove that $\mathbf{d}(s) = \mathbf{d}(t)$, $\mathbf{r}(s) = \mathbf{r}(t)$. By choosing our idempotent to be ss^{-1} we get that $ss^{-1} \leq tt^{-1}$. By symmetry we deduce that $\mathbf{r}(s) = \mathbf{r}(t)$. The fact that $\mathbf{d}(s) = \mathbf{d}(t)$ follows from the same argument using the fact that $(s^{-1}, t^{-1}) \in \mu'$. We have shown that $\mu' \subseteq \mu$.

To prove the converse, suppose that $(s, t) \in \mu$. Let e be an arbitrary idempotent. Then $s^{-1}s = t^{-1}t$ and so $s^{-1}se = t^{-1}te$. Thus $s(s^{-1}se)s^{-1} = t(t^{-1}te)t^{-1}$, which simplifies to $ses^{-1} = tet^{-1}$. It follows that $(s, t) \in \mu'$, as required. \square

We have defined idempotent-separating homomorphisms and we may likewise define idempotent-separating congruences.

Lemma 4.4. *μ is the largest idempotent-separating congruence on S .*

Proof. Let ρ be any idempotent separating-congruence on S and let $(s, t) \in \rho$. Let e be any idempotent. Then $(ses^{-1}, tet^{-1}) \in \rho$ but ρ is idempotent separating and so $ses^{-1} = tet^{-1}$. It follows that $(s, t) \in \mu$. Thus we have shown that $\rho \subseteq \mu$. \square

An inverse semigroup is said to be *fundamental* if μ is the equality relation.

Lemma 4.5. *Let S be an inverse semigroup. Then S/μ is fundamental.*

Proof. Suppose that $\mu(s)$ and $\mu(t)$ are μ -related in S/μ . Every idempotent in S/μ is of the form $\mu(e)$ where $e \in \mathbf{E}(S)$. Thus

$$\mu(s)\mu(e)\mu(s)^{-1} = \mu(t)\mu(e)\mu(t)^{-1}$$

so that $\mu(ses^{-1}) = \mu(tet^{-1})$. But both ses^{-1} and tet^{-1} are idempotents, so that $ses^{-1} = tet^{-1}$ for every $e \in \mathbf{E}(S)$. Thus $(s, t) \in \mu$. \square

Theorem 4.6. *Let S be an inverse semigroup. Then S is fundamental if and only if S is isomorphic to a wide inverse subsemigroup of the Munn semigroup $T_{\mathbf{E}(S)}$.*

Proof. Let S be a fundamental inverse semigroup. By Theorem 4.2, there is a homomorphism $\delta: S \rightarrow T_{\mathbf{E}(S)}$ such that $\ker \delta = \mu$. By assumption, μ is the equality

congruence, and so δ is an injective homomorphism. Thus S is isomorphic to its image in $T_{\mathbf{E}(S)}$, which is a wide inverse subsemigroup.

Conversely, let S be a wide inverse subsemigroup of a Munn semigroup T_E . Clearly, we can assume that $E = \mathbf{E}(S)$. We calculate the maximum idempotent-separating congruence of S . Let $\alpha, \beta \in S$ and suppose that $(\alpha, \beta) \in \mu$ in S . Then $\text{dom } \alpha = \text{dom } \beta$. Let $e \in \text{dom } \alpha$. Then $1_{[e]} \in S$, since S is a wide inverse subsemigroup of $T_{\mathbf{E}(S)}$. By assumption $\alpha 1_{[e]} \alpha^{-1} = \beta 1_{[e]} \beta^{-1}$. It is easy to check that $1_{[\alpha(e)]} = \alpha 1_{[e]} \alpha^{-1}$ and $1_{[\beta(e)]} = \beta 1_{[e]} \beta^{-1}$. Thus $\alpha(e) = \beta(e)$. Hence $\alpha = \beta$, and so S is fundamental. \square

In group theory, congruences are handled using normal subgroups, and in ring theory by ideals. In general semigroup theory, there are no such substructures and so congruences have to be studied in their own right something that is common to most of universal algebra. Even in the case of inverse semigroups, congruences have to be used. However, idempotent-separating homomorphisms are determined by analogues of normal subgroups.

Let $\theta: S \rightarrow T$ be a homomorphism of inverse semigroups. The *Kernel* of θ is defined to be the set K of all elements of S that map to idempotents under θ . Observe that K is a wide inverse subsemigroup of S and it is *self-conjugate* in the sense that $s^{-1}Ks \subseteq K$ for all $s \in S$. We say that K is a *normal inverse subsemigroup* of S .

Remark 4.7. This typographical distinction between *kernels* which are congruences and *Kernels* which are substructures is not entirely happy but convenient for the purposes of this section.

If θ is idempotent-separating then its Kernel satisfies an additional property. If $a \in K$ and if e is any idempotent then $ae = ea$. This motivates the following definition.

For every inverse semigroup S , we define $Z(\mathbf{E}(S))$, the *centralizer of the idempotents*, to be set of all elements of S which commute with every idempotent. The centralizer is a normal inverse subsemigroup and is Clifford. Thus the Kernels of idempotent-separating homomorphisms from S are subsets of the centralizer of the idempotents of S . We now prove that idempotent-separating homomorphisms are determined by their Kernels.

Theorem 4.8. *Let S be an inverse semigroup. Let K be a normal inverse subsemigroup of S contained in $Z(\mathbf{E}(S))$. Define the relation ρ_K by*

$$(s, t) \in \rho_K \Leftrightarrow st^{-1} \in K \text{ and } \mathbf{d}(s) = \mathbf{d}(t).$$

Then ρ_K is an idempotent-separating congruence whose associated Kernel is K .

Proof. We show first that ρ_K is an equivalence relation. Reflexivity and symmetry hold because K is a wide inverse subsemigroup of S . To prove transitivity suppose that $(a, b), (b, c) \in \rho_K$. Then $ab^{-1}, bc^{-1} \in K$ and $\mathbf{d}(a) = \mathbf{d}(b) = \mathbf{d}(c)$. Observe that $ab^{-1}bc^{-1} = ac^{-1} \in K$ and $\mathbf{d}(a) = \mathbf{d}(c)$. Hence $(a, c) \in \rho_K$. Next we show that ρ_K is a congruence. Let $(a, b) \in \rho_K$ and $c \in S$. By assumption, $ab^{-1} \in K$ and $\mathbf{d}(a) = \mathbf{d}(b)$. We prove first that ρ_K is a right congruence by showing that $(ac, bc) \in \rho_K$. Observe that $ac(bc)^{-1} = acc^{-1}b^{-1}$. We may move the idempotent cc^{-1} through b^{-1} by Lemma 2.6. Thus by the fact that K is a wide inverse subsemigroup we have show that $ac(bc)^{-1} \in K$. A simple calculation shows that $\mathbf{d}(ac) = \mathbf{d}(bc)$. We

prove now that ρ_K is a left congruence by showing that $(ca, cb) \in \rho_K$. Observe that $ca(cb)^{-1} = c(ab^{-1})c^{-1}$, but $ab^{-1} \in K$ and K is self-conjugate so that $ca(cb)^{-1} \in K$.

It remains to show that the elements

$$(ca)^{-1}ca = a^{-1}c^{-1}ca \text{ and } (cb)^{-1}cb = b^{-1}c^{-1}cb$$

are equal. Put $e = c^{-1}c$. We shall show that $a^{-1}ea = b^{-1}eb$. Write

$$a^{-1}ea = (a^{-1}ea)(a^{-1}a)(a^{-1}ea).$$

But $a^{-1}a = b^{-1}b$ and so

$$a^{-1}ea = (a^{-1}ea)(b^{-1}b)(a^{-1}ea).$$

Now

$$(a^{-1}ea)(b^{-1}b)(a^{-1}ea) = (a^{-1}e)(ab^{-1})(ab^{-1})^{-1}(ea).$$

But $ab^{-1} \in K$, and K is contained in the centralizer of the idempotents, and so

$$ab^{-1}(ab^{-1})^{-1} = (ab^{-1})^{-1}ab^{-1}.$$

Thus

$$(a^{-1}e)(ab^{-1})(ab^{-1})^{-1}(ea) = (a^{-1}e)(ab^{-1})^{-1}(ab^{-1})(ea),$$

and so

$$a^{-1}ea = (a^{-1}e)(ba^{-1}ab^{-1})(ea).$$

Now

$$(a^{-1}e)(ba^{-1}ab^{-1})(ea) = a^{-1}(ab^{-1}e)^{-1}(ab^{-1}e)a.$$

But $ab^{-1} \in K$, and K is a wide subsemigroup, so that $ab^{-1}e \in K$. Thus because K is contained in the centralizer of the idempotents we have that

$$a^{-1}(ab^{-1}e)^{-1}(ab^{-1}e)a = a^{-1}(ab^{-1}e)(ab^{-1}e)^{-1}a.$$

Thus

$$a^{-1}ea = a^{-1}(ab^{-1}e)(ab^{-1}e)^{-1}a.$$

But $a^{-1}(ab^{-1}e)(ab^{-1}e)^{-1}a = a^{-1}ab^{-1}eb$, so that we in fact have

$$a^{-1}ea = a^{-1}ab^{-1}eb.$$

But then from $a^{-1}a = b^{-1}b$ we obtain $a^{-1}ea = b^{-1}eb$ as required.

We now calculate the Kernel of ρ_K . Let a be in the Kernel of ρ_K . Then there is an idempotent $e \in S$ such that $(a, e) \in \rho_K$. But then $ae \in K$ and $a^{-1}a = e$. Thus $a \in K$. It follows that the Kernel of ρ_K is contained in K . To prove the reverse inclusion, suppose that $a \in K$. Then $a(a^{-1}a) \in K$ and $a^{-1}a = a^{-1}a$. Thus $(a, a^{-1}a) \in \rho_K$. Hence a belongs to the Kernel of ρ_K . \square

The following now confirms what we already suspect.

Proposition 4.9. *Let S be an inverse semigroup. The idempotent-separating congruence determined by $Z(E(S))$ is μ .*

Proof. We calculate the Kernel of μ . Suppose that $s\mu e$ where e is an idempotent. Let f be an arbitrary idempotent. Then $sfs^{-1}\mu ef$ and $fss^{-1}\mu ef$. Thus $sfs^{-1}\mu fss^{-1}$ and so $sfs^{-1} = fss^{-1}$. It follows that $sf = fs$ and $s \in Z(E(S))$. Conversely, let $s \in Z(E(S))$. Then $s\mu ss^{-1}$. \square

The following result provides a useful criterion for a semigroup to be fundamental.

Proposition 4.10. *Let S be an inverse semigroup. Then S is fundamental if and only if $Z(\mathbf{E}(S)) = \mathbf{E}(S)$.*

Proof. Suppose that S is fundamental. Let $a \in Z(\mathbf{E}(S))$. By Proposition 4.9, $\text{Ker } \mu = Z(\mathbf{E}(S))$. Thus $(a, e) \in \mu$ for some $e \in \mathbf{E}(S)$. But then $a = e$, since μ is equality, and so a is an idempotent. Thus $Z(\mathbf{E}(S)) = \mathbf{E}(S)$.

Conversely, suppose that $Z(\mathbf{E}(S)) = \mathbf{E}(S)$. Let $(a, b) \in \mu$. Then

$$(ab^{-1}, bb^{-1}) \in \mu,$$

and so $ab^{-1} \in \text{Ker } \mu$. But $\text{Ker } \mu = Z(\mathbf{E}(S))$ by Proposition 4.9, and so $ab^{-1} \in Z(\mathbf{E}(S))$. Thus ab^{-1} is an idempotent, by assumption. But then $ab^{-1} = bb^{-1}$ since μ is idempotent-separating, which gives $ab^{-1}b = b$. But $\mathbf{d}(a) = \mathbf{d}(b)$ and so $a = b$. \square

A topological space X is said to be T_0 if for each pair of elements $x, y \in X$ there exists an open set which contains one but not both of x and y . A *base* for a topological space is a set of open sets β such that every open set of the topology is a union of elements of β . Let X be an arbitrary set and β a set of subsets of X whose union is X and with the property that the intersection of any two elements of β is a union of elements of β . Then a topology can be defined on X by defining the open sets to be the unions of elements of β .

As in Example 2.9, the inverse semigroup of all homeomorphisms between open subsets of X is denoted by $\Gamma(X)$. An inverse subsemigroup S of $\Gamma(X)$ is said to be *topologically complete* if the set-theoretic domains of the elements of S form a base for the topology.

Theorem 4.11. *An inverse semigroup is fundamental if and only if it is isomorphic to a topologically complete inverse semigroup on a T_0 -space.*

Proof. Let S be a fundamental inverse semigroup. We can assume by Theorem 4.6, that S is a wide inverse subsemigroup of a Munn semigroup T_E . Put $\beta = \{e^\perp : e \in E\}$. Clearly, E is the union of the elements of β , and β is closed under finite intersections. Thus β is the base of a topology on the set E . With respect to this topology, each element of S is a homeomorphism between open subsets of E . It remains to show that this topology is T_0 . Let $e, f \in E$ be distinct idempotents. If $f \leq e$ then f^\perp is an open set containing f but not e . If $f \not\leq e$ then e^\perp is an open set containing e but not f . Thus the topology is T_0 .

Conversely, let S be a topologically complete inverse subsemigroup of the inverse semigroup $\Gamma(X) = I(X, \tau)$ where the topology τ is T_0 and $\beta = \{\text{dom } \alpha : \alpha \in S\}$ is a base for τ . We shall prove that S is fundamental by showing that the centralizer of the idempotents of S contains only idempotents (Proposition 4.10). Let $\phi \in S \setminus \mathbf{E}(S)$. Then there exists $x \in \text{dom } \phi$ such that $\phi(x) \neq x$, because ϕ is not an idempotent. Since τ is T_0 , there exists an open set U such that

$$\text{either } (\phi(x) \in U \text{ and } x \notin U) \text{ or } (\phi(x) \notin U \text{ and } x \in U).$$

Since β is a basis for the topology, $U = \bigcup B_i$ for some $B_i \in \beta$. It follows that there is a $B = B_i \in \beta$ such that

$$\text{either } (\phi(x) \in B \text{ and } x \notin B) \text{ or } (\phi(x) \notin B \text{ and } x \in B).$$

Observe that $1_B \in S$ since $B = \text{dom } \alpha$ for some $\alpha \in S$. Thus the elements $\phi 1_B$ and $1_B \phi$ belong to S . In the first case, $\phi(x) \in B$ and $x \notin B$, so that whereas $(\phi 1_B)(x)$

is not defined, $(1_B\phi)(x)$ is defined. Thus $\phi \notin Z(E(S))$. In the second case, $(\phi 1_B)(x)$ is defined and $(1_B\phi)(x)$ is not defined. Thus once again $\phi \notin Z(E(S))$. Hence in either case $\phi \notin Z(E(S))$. \square

Let S be an arbitrary inverse semigroup, let its image under the Munn representation be T , and let K be the centralizer of the idempotents of S . Then S is an extension of K by T where the former is a presheaf of groups and the latter is a pseudogroup of transformations.

Theorem 4.12. *Every inverse semigroup is an idempotent-separating extension of a presheaf of groups by a pseudogroup of transformations.*

4.2. Congruence-free inverse semigroups. A useful application of fundamental inverse semigroups is in characterizing those semigroups which are congruence-free. I shall concentrate only on the case of inverse semigroups with zero. Douglas Munn once remarked to me that this was one of the few instances where the theory for inverse semigroups with zero was easier than it was for the one without. We shall need a sequence of definitions before we can state our main result.

Although ideals are useful in semigroup theory, the connection between ideals and congruences is weaker for semigroups than it is for rings. If ρ is a congruence on a semigroup with zero S , then the set $I = \rho(0)$ is an ideal of S ; however, examples show that the congruence is not determined by this ideal. Nevertheless, ideals can be used to construct some congruences on semigroups. Let I be an ideal in the semigroup S . Define a relation ρ_I on S by:

$$(s, t) \in \rho_I \Leftrightarrow \text{either } s, t \in I \text{ or } s = t.$$

Then ρ_I is a congruence. The quotient semigroup S/ρ_I is isomorphic to the set $S \setminus I \cup \{0\}$ (we may assume that $0 \notin S \setminus I$) equipped with the following product: if $s, t \in S \setminus I$ then their product is st if $st \in S \setminus I$, all other products are defined to be 0. Such quotients are called *Rees quotients*.

There is also a way of constructing congruences from subsets. Let S be a semigroup and let $L \subseteq S$. Define a relation ρ_L on S by:

$$(s, t) \in \rho_L \Leftrightarrow (\forall a, b \in S)(asb \in L \Leftrightarrow atb \in L).$$

Then ρ_L is a congruence on S , called the *syntactic congruence* of L .

An inverse semigroup with zero S is said to be *0-simple* if it contains at least one non-zero element and the only ideals are $\{0\}$ and S . An inverse semigroup is said to be *congruence-free* if its only congruences are equality and the universal congruence. Thus congruence-free-ness is much stronger than 0-simplicity. A congruence ρ is said to be *0-restricted* if the ρ -class containing 0 is just 0. Finally, define ξ to be the syntactic congruence of the subset $\{0\}$.

Lemma 4.13. *The congruence ξ is the maximum 0-restricted congruence.*

Proof. Let ρ be a 0-restricted congruence on S and let spt . Suppose that $asb = 0$. But $asb\xi atb$ and so since ρ is 0-restricted, we have that $atb = 0$. By symmetry we deduce that $a\xi b$. Thus $\rho \subseteq \xi$, as required. \square

Lemma 4.14. *Let S be an inverse semigroup with zero.*

- (1) $\mu \subseteq \xi$.
- (2) *The congruence ξ restricted to $E(S)$ is the syntactic congruence determined by zero on $E(S)$.*

Proof. (1) Let $s\mu t$. Suppose that $asb = 0$ then $asb\mu atb$ and so $atb = 0$. By symmetry this shows that $s\xi t$.

(2) Let e and f be idempotents. Suppose that for all idempotents i we have that $ie = 0$ iff $if = 0$. Let $aeb = 0$. Then $a^{-1}aebb^{-1} = 0$. Thus $a^{-1}abb^{-1}e = 0$ and so $a^{-1}abb^{-1}f = 0$. Hence $a^{-1}afbb^{-1} = 0$ and so $afb = 0$. The reverse direction is proved similarly. \square

An inverse semigroup with zero is said to be *0-disjunctive* if ξ is the equality relation.

Proposition 4.15. *An inverse semigroup S is 0-disjunctive if and only if $E(S)$ is 0-disjunctive and S is fundamental.*

Proof. If S is 0-disjunctive it follows by Lemma 4.14 that $E(S)$ is 0-disjunctive and S is fundamental. Suppose that $E(S)$ is 0-disjunctive and S is fundamental. Then ξ restricted to $E(S)$ is the equality relation and so ξ is idempotent-separating. Thus by Lemma 4.4 $\xi \subseteq \mu$. But S is fundamental and so μ is the equality relation and so ξ is the equality relation. \square

Lemma 4.16. *Let E be a meet semilattice with zero. Then the following are equivalent.*

- (1) E is 0-disjunctive.
- (2) For all distinct $e, f \in E$ nonzero there exists $g \in E$ such that either $e \wedge g \neq 0$ and $f \wedge g = 0$ or $e \wedge g = 0$ and $f \wedge g \neq 0$.
- (3) For all $0 \neq f < e$ there exists $0 \neq g \leq e$ such that $f \wedge g = 0$.

Proof. (1) \Rightarrow (2). This is immediate from the definition.

(2) \Rightarrow (3). Let $0 \neq f < e$. Then there exists g' such that $g' \wedge f = 0$ and $g' \wedge e \neq 0$ or $g' \wedge f \neq 0$ and $g' \wedge e = 0$. Clearly the second case cannot occur. Put $g = g' \wedge e$. Then $g \leq e$, $g \neq 0$ and $g \wedge f = 0$, as required.

(3) \Rightarrow (1). Suppose that $e\xi f$ where e and f are both non-zero. Then $e\xi(e \wedge f)$ and so $e \wedge f \neq 0$. Suppose that $e \wedge f \neq e$. Then there exists $0 \neq g \leq e$ such that $(e \wedge f) \wedge g = 0$. But clearly $e \wedge g \neq 0$. We therefore have a contradiction and so $e \wedge f = e$. Similarly $e \wedge f = f$ and so $e = f$, as required. \square

Remark 4.17. Property (3) in the above lemma has a Boolean ‘feel’ to it. If $e < f$ are non-zero then $f\bar{e} < f$ and $e \wedge f\bar{e} = 0$. Thus Boolean algebras are automatically 0-disjunctive. This has implications in the study of Boolean inverse semigroups. See Section 5.

We may now state the characterization of congruence-free inverse semigroups with zero.

Theorem 4.18. *An inverse semigroup with zero S is congruence-free if and only if S is fundamental, 0-simple and $E(S)$ is 0-disjunctive.*

Proof. Suppose that S is congruence-free. Then μ is equality, there are no non-trivial ideals and ξ is equality. Thus S is fundamental, 0-simple and $E(S)$ is 0-disjunctive.

To prove the converse, suppose that S is fundamental, 0-simple and $E(S)$ is 0-disjunctive. Let ρ be a congruence on S which is not the universal relation. Then $\rho(0)$ is an ideal which is not S . Thus it must be equal to $\{0\}$. It follows that ρ is a 0-restricted congruence and so $\rho \subseteq \xi$. But by Proposition 4.15, ξ is the equality congruence and so ρ is the equality congruence. \square

The above theorem will be a useful criterion for congruence-free-ness once we have a nice characterization of 0-simplicity. This involves the one Green's relation we have yet to define. Let S be an inverse semigroup. Define

$$(s, t) \in \mathcal{J} \Leftrightarrow SsS = StS.$$

It is always true that $\mathcal{D} \subseteq \mathcal{J}$. The meaning of the \mathcal{J} -relation for inverse semigroups is clarified by the following result.

Lemma 4.19. *Let S be an inverse semigroup. Then $a \in SbS$ if and only if there exists $u \in S$ such that $a\mathcal{D}u \leq b$.*

Proof. Let $a \in SbS$. Then $a = xby$ for some $x, y \in S$. By Proposition 2.23, there exist elements x', y' and b' such that $a = x' \cdot b' \cdot y'$ is a restricted product where $x' \leq x$, $b' \leq b$ and $a' \leq a$. Hence $a\mathcal{D}b'$ which, together with $b' \leq b$, gives $a\mathcal{D}b' \leq b$. Conversely, suppose that $a\mathcal{D}b' \leq b$. From $a\mathcal{D}b'$ we have that $a\mathcal{J}b'$, and from $b' \leq b$ we have that $Sb'S \subseteq SbS$. Thus $a \in SbS$. \square

Lemma 4.20. *Let S be an inverse semigroup with zero. Then it is 0-simple if and only if $S \neq \{0\}$ and the only \mathcal{J} -classes are $\{0\}$ and $S \setminus \{0\}$.*

Proof. Let S be 0-simple and let $s, t \in S$ be a pair of non-zero elements. Both SsS and StS are ideals of S and so must be equal. Thus $(s, t) \in \mathcal{J}$. Conversely, suppose that the only non-zero \mathcal{J} -class is $S \setminus \{0\}$. Let I be any non-zero ideal of S . Let $s \in I$ and $t \in S$ be non-zero elements. By assumption, $(s, t) \in \mathcal{J}$. Thus $t = asb$ for some $a, b \in S$ and so $t \in I$. Hence $I = S \setminus \{0\}$. \square

Proposition 4.21. *Let S be an inverse semigroup with zero.*

- (1) *S is 0-simple if and only if for any two non-zero elements s and t in S there exists an element s' such that $s\mathcal{D}s' \leq t$.*
- (2) *S is 0-simple if and only if for any two non-zero idempotents e and f in S there exists an idempotent i such that $e\mathcal{D}i \leq f$.*

Proof. (1) By Lemma 4.19, an inverse semigroup is 0-simple if it consists of exactly two \mathcal{J} -class $\{0\}$ and $S \setminus \{0\}$. Thus any two non-zero elements of S are \mathcal{J} -related. The result is now immediate by Lemma 4.18.

(2) Suppose the condition on the idempotents holds. Let $s, t \in S$ be a pair of non-zero elements. Then $e = ss^{-1}$ and $f = tt^{-1}$ are non-zero idempotents and so, by assumption, there is an idempotent i such that $e\mathcal{D}i \leq f$. Put $u = it$. Then $u \leq t$, and $uu^{-1} = it(it)^{-1} = itt^{-1} = if = i$. Thus $s\mathcal{D}u \leq t$. The proof of the converse is straightforward. \square

5. BOOLEAN INVERSE MONOIDS

Let S be an inverse monoid with zero. We say that it is *distributive* if it satisfies the following two conditions:

- (1) Compatible pairs of elements have a join. We shall call these *compatible joins*.
- (2) Multiplication distributes over compatible joins.

The semilattice of idempotents of a distributive inverse monoid is a distributive lattice (with both top and bottom). A distributive inverse monoid is said to be *Boolean* if its semilattice of idempotents is actually a Boolean algebra. Our perspective is that Boolean inverse monoids are non-commutative generalizations of Boolean algebras.

Examples 5.1. Here are a list of some examples of Boolean inverse monoids.

- (1) Symmetric inverse monoids $I(X)$ are Boolean inverse monoids. Its Boolean algebra of idempotents is isomorphic to the power set Boolean algebra $P(X)$. If X is finite with n elements then we denote the symmetric inverse monoid on an n -element set by I_n .
- (2) Groups with zero adjoined. We denote these by G^0 . These may look like chimeras but they are honest-to-goodness Boolean inverse monoids whose Boolean algebra of idempotents is isomorphic to the 2-element Boolean algebra.
- (3) R_n the set of all $n \times n$ rook matrices [29]. These are all $n \times n$ matrices over the numbers 0 and 1 such that each row and each column contains at most one non-zero entry. In fact, R_n is isomorphic to I_n .
- (4) $R_n(G^0)$ the set of all $n \times n$ rook matrices over a group with zero [23]. These are all $n \times n$ matrices over the group with zero G^0 in which each row and each column contains at most one non-zero entry.

REFERENCES

- [1] A. Coudron, Sur les extensions de demi-groupe réciproques, *Bull. Soc. Roy. Sci. Liège*, **37** (1968), 409–419.
- [2] H. D’Alarcao, Idempotent separating extensions of inverse semigroups, *J. Austral. Math. Soc.* **9** (1969), 211–217.
- [3] Ch. Ehresmann, *Oeuvres complètes et commentées*, (ed A. C. Ehresmann) Supplements to *Cahiers de topologie et géométrie différentielle*, Amiens, 1980–83.
- [4] R. Exel, Inverse semigroups and combinatorial C^* -algebras, *Bull. Braz. Math. Soc. (N.S.)* **39** (2008), 191–313.
- [5] J.-Y. Girard, The geometry of interaction III: accommodating the additives, in *Advances in linear logic* (eds J.-Y. Girard, Y. Lafont, L. Regnier) Cambridge University Press, 1995.
- [6] J. M. Howie, *Fundamentals of semigroup theory*, Clarendon Press, Oxford, 1995.
- [7] D. G. Jones, M. V. Lawson, Graph inverse semigroups: their characterization and completion, *J. Alg.* **409** (2014), 444–473.
- [8] P. T. Johnstone, *Stone spaces*, CUP, 1982.
- [9] J. Kellendonk, The local structure of tilings and their integer group of coinvariants, *Commun. Math. Phys.* **187** (1997), 115–157.
- [10] G. Kudryavtseva, M. V. Lawson, D. H. Lenz, P. Resende, Invariant means on Boolean inverse monoids, *Semigroup Forum* **92** (2016), 77–101.
- [11] A. Kumjian, On localization and simple C^* -algebras, *Pacific J. Maths* **112** (1984), 141–192.
- [12] M. V. Lawson, *Inverse semigroups*, the theory of partial symmetries, World Scientific, 1998.
- [13] M. V. Lawson, The polycyclic monoids P_n and the Thompson groups $V_{n,1}$, *Comm. Algebra* **35** (2007), 4068–4087.
- [14] M. V. Lawson, A non-commutative generalization of Stone duality, *J. Aust. Math. Soc.* **88** (2010), 385–404.
- [15] M. V. Lawson, Compactable semilattices, *Semigroup Forum* **81** (2010), 187–199.
- [16] M. V. Lawson, Non-commutative Stone duality: inverse semigroups, topological groupoids and C^* -algebras, *IJAC* **22**, 1250058 (2012) DOI:10.1142/S0218196712500580.
- [17] M. V. Lawson, D. H. Lenz, Pseudogroups and their étale groupoids, *Adv. Math.* **244** (2013), 117–170.
- [18] M. V. Lawson, P. Scott, AF inverse monoids and the structure of countable MV-algebras, *J. Pure Appl. Alg.* **221** (2017), 45–74.
- [19] M. V. Lawson, S. Margolis, B. Steinberg, The étale groupoid of an inverse semigroup as a groupoid of filters, *J. Austral. Math. Soc.* **94** (2014), 234–256.
- [20] H. Lausch, Cohomology of inverse semigroups, *J. Algebra* **35** (1975), 273–303.
- [21] M. Loganathan, Cohomology of inverse semigroups, *J. Algebra* **70** (1981), 375–393.
- [22] S. Mac Lane, *Categories for the working mathematician*, Springer-Verlag, Berlin, 1971.

- [23] M. E. Malandro, Fast Fourier transforms for finite inverse semigroups, *J. Algebra* **324** (2010), 282–312.
- [24] A. L. T. Paterson, *Groupoids, inverse semigroups, and their operator algebras*, Progress in Mathematics, **170**, Birkhäuser, Boston, 1998.
- [25] M. Petrich, *Inverse semigroups*, John Wiley & Sons, 1984.
- [26] J. Renault, *A groupoid approach to C^* -algebras*, Springer-Verlag, 1980.
- [27] P. Resende, Lectures on étale groupoids, inverse semigroups and quantales, Lecture Notes, August, 2006.
- [28] P. Resende, Etale groupoids and their quantales, *Adv. Math.* **208** (2007), 147–209.
- [29] L. Solomon, Representations of rook monoids, *J. Algebra* **256** (2002), 309–342.
- [30] A. R. Wallis, *Semigroups and category-theoretic approaches to partial symmetry*, PhD Thesis, Heriot-Watt University, Edinburgh, 2013.
- [31] F. Wehrung, Refinement monoids, equidivisibility types, and Boolean inverse monoids, *Lecture Notes in Mathematics* 2188, Springer, 2017.
- [32] F. Wehrung, Varieties of Boolean inverse semigroups, arXiv arXiv:1610.07447.

DEPARTMENT OF MATHEMATICS AND THE MAXWELL INSTITUTE FOR MATHEMATICAL SCIENCES,
HERIOT-WATT UNIVERSITY, RICcarton, EDINBURGH EH14 4AS, UNITED KINGDOM